

REPORT DOCUMENTATION PAGE*Form Approved*
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 24-26 October 1995	3. REPORT TYPE AND DATES COVERED Proceedings, 1995	
4. TITLE AND SUBTITLE Global Statistics on Cloud Optical Depths From Satellite and Lidar Observations			5. FUNDING NUMBERS N/A	
6. AUTHOR(S) Donald Wylie				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Space Science and Engineering Center, University of Wisconsin-Madison, Madison, Wisconsin			8. PERFORMING ORGANIZATION REPORT NUMBER N/A	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) SERDP 901 North Stuart St. Suite 303 Arlington, VA 22203			10. SPONSORING / MONITORING AGENCY REPORT NUMBER N/A	
11. SUPPLEMENTARY NOTES Presented at Cloud Impacts on DOD Operations and Systems 1995 Conference, U.S. Air Force Phillips Laboratory-Science Center, Hanscom Air Force Base, Massachusetts, 24-26 October 1995. No copyright is asserted in the United States under Title 17, U.S. code. The U.S. Government has a royalty-free license to exercise all rights under the copyright claimed herein for Government purposes. All other rights are reserved by the copyright owner.				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release: distribution is unlimited				12b. DISTRIBUTION CODE A
13. ABSTRACT (Maximum 200 Words) Six years of cloud data have been collected globally from the NOAA satellite series. The High Resolution Infrared Radiometer Spectrometer (HIRS) data were used to detect clouds and estimate their optical depths in the 11 micron infrared window. The cloud detection algorithm has been called the CO ₂ Slicing Algorithm because it uses the 13-15 micron infrared channels where partial CO ₂ absorption occurs, to detect partially transparent clouds and correctly determine their altitude. This algorithm is designed to be sensitive to upper tropospheric cirrus clouds which are difficult to detect. The frequency of these clouds along with their global distribution and seasonal changes have been reported at past CIDOS conferences.				
14. SUBJECT TERMS NOAA satellite series, HIRS, lidar observation, tropospheric cirrus clouds, SERDP				15. NUMBER OF PAGES 1
				16. PRICE CODE N/A
17. SECURITY CLASSIFICATION OF REPORT unclass	18. SECURITY CLASSIFICATION OF THIS PAGE unclass	19. SECURITY CLASSIFICATION OF ABSTRACT unclass	20. LIMITATION OF ABSTRACT UL	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)
Prescribed by ANSI Std. Z39-18
298-102

CLOUD IMPACTS ON DOD OPERATIONS AND SYSTEMS 1995 CONFERENCE

U.S. Air Force Phillips Laboratory - Science Center
Hanscom Air Force Base, Massachusetts
24-26 October 1995

19980806 036

CIDOS - 95

Cloud Modeling and Data for Defense Simulation Activities
"Emphasizing Sufficient Physical Reality in Simulating Clouds"

PL-TR-95-2129

Environmental Research Papers, No. 1179


**PREPRINT OF THE CLOUD IMPACTS ON
DoD OPERATIONS AND SYSTEMS
1995 CONFERENCE (CIDOS-95)**

Editor

Donald D. Grantham

1 October 1995

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED



**PHILLIPS LABORATORY
Directorate of Geophysics
Air Force Materiel Command
Hanscom Air Force Base, MA 01731-3010**

GLOBAL STATISTICS ON CLOUD OPTICAL DEPTHS FROM SATELLITE AND LIDAR OBSERVATIONS

Donald Wylie

Space Science and Engineering Center
University of Wisconsin-Madison
Madison, Wisconsin

Six years of cloud data have been collected globally from the NOAA satellite series. The High Resolution Infrared Radiometer Spectrometer (HIRS) data were used to detect clouds and estimate their optical depths in the 11 micron infrared window. The cloud detection algorithm has been called the CO2 Slicing Algorithm because it uses the 13-15 micron infrared channels where partial CO2 absorption occurs, to detect partially transparent clouds and correctly determine their altitude. This algorithm is designed to be sensitive to upper tropospheric cirrus clouds which are difficult to detect. The frequency of these clouds along with their global distribution and seasonal changes have been reported at past CIDOS conferences.

The optical depths of semi-transparent cirrus clouds from the CO2 Slicing data. A detailed comparison of infrared optical depths derived from satellite data to optical depths measured by the University of Wisconsin's High Spectral Resolution Lidar (HSRL) has been made. The HSRL is discussed in a separate paper by E. Eloranta in these proceedings. Coincident satellite and lidar optical depth data were collected on 21 days in the past two years. The optical depth comparison exhibited scatter because of differences in which each sensor scanned the clouds. The satellite scanned over a large area with a lower resolution field of view; typically 1-8 km in diameter, while the lidar scanned a narrow line through the cloud < 1 m wide from the wind advection of the cloud over the lidar. The spatial variability of cloud density produced most of the disagreement between the two measurements. The mean optical depths, however, did agree after accounting for differences in the radiative physics of scattering and absorption between the visible and infrared measurements. A satellite bias toward larger optical depths was found for very thin cirrus clouds at the minimum detectable density of the satellite system. This bias occurred because very thin clouds had to be at least 2 K colder than surrounding cloud free backgrounds to be distinguished from noise and natural scene variability on the satellite images. This constraint imposed a minimum IR optical depth of 0.05 (0.10 in the visible) which is near the "invisible cirrus" category.

Thicker clouds appeared to be correctly measured by the satellite in spite of the fact that they are often 3-5 km thick and composed of multiple layers.

Manuscript not available at time of printing. Please contact author for information.